





What is claimed is:

1. A method for the treatment of hepatitis B virus (HBV) infection comprising administering an effective amount of a compound selected from the group consisting of formulas [I]-[IV]

5 below and mixtures of two or more thereof:



wherein:



E is selected from the group consisting of H, OH, OMe, SH, SMe, NH<sub>2</sub>, NHMe, N<sub>3</sub>, F, Cl, Br, CO<sub>2</sub>H, CO<sub>2</sub>-alkyl, OPh, OPhNO<sub>2</sub>, NO, NO<sub>2</sub>, SCN, OCN, NCS, NCO, SOMe, SO<sub>2</sub>Me;

X is selected from the group consisting of O, S, NH, CH<sub>2</sub>, CHF, CF<sub>2</sub>;

Y is selected from the group consisting of CH<sub>2</sub>, NH, NOH, NMe, NEt, NOME, CHF, CF<sub>2</sub>;

Z is selected from the group consisting of H, OH, OMe, SH, SMe, F, Cl, Br, I, NH<sub>2</sub>, NHMe;

15 B is a base selected from the group consisting of




R<sup>2</sup> is selected from the group consisting of O, S, NH, NR;

R<sup>5</sup> is selected from the group consisting of H, branched or unbranched lower alkyl having from 1-5 carbon atoms, F, Cl, Br, I, CH=CH<sub>2</sub>, CH=CHBr, Ph, Ac, OMe, OPh, NO, NO<sub>2</sub>, NH<sub>2</sub>, NHR;

R<sup>6</sup> and R<sup>8</sup> are the same or different and are independently selected from the group consisting of H, OH, OMe, SH, SMe, F, Cl, Br, I, NH<sub>2</sub>, NHMe, NMe<sub>2</sub>;

R is independently selected from the group consisting of




or a pharmaceutically acceptable salt or prodrug thereof, optionally in combination with a pharmaceutically acceptable carrier.





2. The method of Claim 1, further comprising administering the compound in combination or alternation with one or more additional anti-HBV agents.

3. The method of Claim 2, wherein the additional anti-HBV agent is selected from the group consisting of FTC (the (-)-enantiomer or the racemate), L-FMAU, interferon, beta-D-dioxolanyl-guanine (DXG), beta-D-dioxolanyl-2,6-diaminopurine (DAPD), beta-D-dioxolanyl-6-chloropurine (ACP), beta-D-dioxolanyl-2-aminopurine (ADP), famciclovir, penciclovir, bis-POM PMEA (adefovir dipivoxil); lobucavir, ganciclovir, ribavirin, lamivudine (3TC), L-thymidine (L-dT), L-2'-deoxycytidine (L-dT), L-2'-deoxycytidine-3',5'-di-O-valyl (D or L), entecavir (BMS-200475), adefovir, L-D4FC, D-D4FC, and mycophenolic acid (an IMPDH inhibitor).

4. The method of Claim 1, wherein the compound is selected from the group consisting of



5. A method for the treatment of hepatitis C virus (HCV) infection comprising administering an effective amount of a compound selected from the group consisting of formulas [I]-[IV] below and mixtures of two or more thereof:



wherein:



E is selected from the group consisting of H, OH, OMe, SH, SMe, NH<sub>2</sub>, NHMe, N<sub>3</sub>, F, Cl, Br, CO<sub>2</sub>H, CO<sub>2</sub>-alkyl, OPh, OPhNO<sub>2</sub>, NO, NO<sub>2</sub>, SCN, OCN, NCS, NCO, SMe, SO<sub>2</sub>Me;

X is selected from the group consisting of O, S, NH, CH<sub>2</sub>, CHF, CF<sub>2</sub>;

Y is selected from the group consisting of CH<sub>2</sub>, NH, NOH, NMe, NEt, NOME, CHF, CF<sub>2</sub>;

Z is selected from the group consisting of H, OH, OMe, SH, SMe, F, Cl, Br, I, NH<sub>2</sub>, NHMe;

B is a base selected from the group consisting of




R<sup>2</sup> is selected from the group consisting of O, S, NH, NR;

R<sup>5</sup> is selected from the group consisting of H, branched or unbranched lower alkyl having from 1-5 carbon atoms, F, Cl, Br, I, CH=CH<sub>2</sub>, CH=CHBr, Ph, Ac, OMe, OPh, NO, NO<sub>2</sub>, NH<sub>2</sub>, NHR;

R<sup>6</sup> and R<sup>8</sup> are the same or different and are independently selected from the group consisting of H, OH, OMe, SH, SMe, F, Cl, Br, I, NH<sub>2</sub>, NHMe, NMe<sub>2</sub>;

R is independently selected from the group consisting of




or a pharmaceutically acceptable salt or prodrug thereof, optionally in combination with a pharmaceutically acceptable carrier.





6. The method of Claim 5, further comprising administering the compound in combination or alternation with one or more additional anti-HCV agents.

7. The method of Claim 6, wherein the additional HCV agent is selected from the group consisting of interferon, macrokine, heptazyme, ribavarin (D and L), amantadine, ofloxacin, zadaxin and reticulose.

8. The method of Claim 5, wherein the compound is selected from the group consisting of



9. A method for the treatment of hepatitis D virus (HDV) infection comprising administering an effective amount of a compound selected from the group consisting of formulas [I]-[IV] below and mixtures of two or more thereof:



wherein:



E is selected from the group consisting of H, OH, OMe, SH, SMe, NH<sub>2</sub>, NHMe, N<sub>3</sub>, F, Cl, Br, CO<sub>2</sub>H, CO<sub>2</sub>-alkyl, OPh, OPhNO<sub>2</sub>, NO, NO<sub>2</sub>, SCN, OCN, NCS, NCO, SOMe, SO<sub>2</sub>Me;

X is selected from the group consisting of O, S, NH, CH<sub>2</sub>, CHF, CF<sub>2</sub>;

Y is selected from the group consisting of CH<sub>2</sub>, NH, NOH, NMe, NEt, NOME, CHF, CF<sub>2</sub>;

Z is selected from the group consisting of H, OH, OMe, SH, SMe, F, Cl, Br, I, NH<sub>2</sub>, NHMe;

B is a base selected from the group consisting of




R<sup>2</sup> is selected from the group consisting of O, S, NH, NR;

R<sup>5</sup> is selected from the group consisting of H, branched or unbranched lower alkyl having from 1-5 carbon atoms, F, Cl, Br, I, CH=CH<sub>2</sub>, CH=CHBr, Ph, Ac, OMe, OPh, NO, NO<sub>2</sub>, NH<sub>2</sub>, NHR;

- 5 R<sup>6</sup> and R<sup>8</sup> are the same or different and are independently selected from the group consisting of H, OH, OMe, SH, SMe, F, Cl, Br, I, NH<sub>2</sub>, NHMe, NMe<sub>2</sub>;

R is independently selected from the group consisting of




or a pharmaceutically acceptable salt or prodrug thereof, optionally in combination with a pharmaceutically acceptable carrier.





- 10 10. The method of Claim 9, further comprising administering the compound in combination or alternation with one or more additional anti-HDV agents.
11. The method of Claim 10, wherein the additional HDV agent is selected from the group consisting of FTC (the (-)-enantiomer or the racemate), L-FMAU, interferon, beta-D-dioxolanyl-guanine (DXG), beta-D-dioxolanyl-2,6-diaminopurine (DAPD), beta-D-dioxolanyl-6-chloropurine (ACP), beta-D-dioxolanyl-2-aminopurine (ADP), famciclovir, penciclovir, bis-POM PMEA (adefovir dipivoxil); lobucavir, ganciclovir, ribavirin, lamivudine (3TC), L-thymidine (L-dT), L-2'-deoxycytidine (L-dT), L-2'-deoxycytidine-3',5'-di-O-valyl (D or L), entecavir (BMS-200475), adefovir, L-D4FC, D-D4FC, and mycophenolic acid (an IMPDH inhibitor).
- 15



12. The method of Claim 9, wherein the compound is selected from the group consisting of



13. A method for the treatment of HIV or AIDS comprising administering an effective amount of a compound selected from the group consisting of formulas [I]-[IV] below and mixtures of two or more thereof:



wherein:



E is selected from the group consisting of H, OH, OMe, SH, SMe, NH<sub>2</sub>, NHMe, N<sub>3</sub>, F, Cl, Br, CO<sub>2</sub>H, CO<sub>2</sub>-alkyl, OPh, OPhNO<sub>2</sub>, NO, NO<sub>2</sub>, SCN, OCN, NCS, NCO, SOMe, SO<sub>2</sub>Me;

X is selected from the group consisting of O, S, NH, CH<sub>2</sub>, CHF, CF<sub>2</sub>;

Y is selected from the group consisting of CH<sub>2</sub>, NH, NOH, NMe, NEt, NOME, CHF, CF<sub>2</sub>;

Z is selected from the group consisting of H, OH, OMe, SH, SMe, F, Cl, Br, I, NH<sub>2</sub>, NHMe;

B is a base selected from the group consisting of




R<sup>2</sup> is selected from the group consisting of O, S, NH, NR;

[illegible]

15


turbo







15. The method of Claim 14, wherein the additional anti-HIV agent is selected from the group consisting of (-) or racemic 2-hydroxymethyl-5-(5-fluorocytosin-1-yl)-1,3-oxathiolane (FTC); the (-)-enantiomer of 2-hydroxymethyl-5-(cyostin-1-yl)-1,3-oxathiolane (3TC);

0713672.01

16. The method of claim 13, wherein the compound is selected from the group consisting of



17. A pharmaceutical composition for the treatment of HBV comprising a combination of an effective amount of an anti-HBV agent and an effective amount of a compound selected from the group consisting of formulas [I]-[IV] below and mixtures of two or more thereof:



wherein:



E is selected from the group consisting of H, OH, OMe, SH, SMe, NH<sub>2</sub>, NHMe, N<sub>3</sub>, F, Cl, Br, CO<sub>2</sub>H, CO<sub>2</sub>-alkyl, OPh, OPhNO<sub>2</sub>, NO, NO<sub>2</sub>, SCN, OCN, NCS, NCO, SMe, SO<sub>2</sub>Me;

X is selected from the group consisting of O, S, NH, CH<sub>2</sub>, CHF, CF<sub>2</sub>;

Y is selected from the group consisting of CH<sub>2</sub>, NH, NOH, NMe, NEt, NMe, CHF, CF<sub>2</sub>;

Z is selected from the group consisting of H, OH, OMe, SH, SMe, F, Cl, Br, I, NH<sub>2</sub>, NHMe;

B is a base selected from the group consisting of




R<sup>2</sup> is selected from the group consisting of O, S, NH, NR;

R<sup>5</sup> is selected from the group consisting of H, branched or unbranched lower alkyl having from 1-5 carbon atoms, F, Cl, Br, I, CH=CH<sub>2</sub>, CH=CHBr, Ph, Ac, OMe, OPh, NO, NO<sub>2</sub>, NH<sub>2</sub>, NHR;


R<sup>6</sup> and R<sup>8</sup> are the same or different and are independently selected from the group consisting of H, OH, OMe, SH, SMe, F, Cl, Br, I, NH<sub>2</sub>, NHMe, NMe<sub>2</sub>;

R is independently selected from the group consisting of







or a pharmaceutically acceptable salt or prodrug thereof, in combination with a pharmaceutically acceptable carrier.

18. The composition of Claim 17, wherein the compound is selected from the group consisting of



19. A pharmaceutical composition for the treatment of HCV comprising an anti-HCV agent and an effective amount of a compound selected from the group consisting of formulas [I]-[IV] below and mixtures of two or more thereof:



wherein:



E is selected from the group consisting of H, OH, OMe, SH, SMe, NH<sub>2</sub>, NHMe, N<sub>3</sub>, F, Cl, Br, CO<sub>2</sub>H, CO<sub>2</sub>-alkyl, OPh, OPhNO<sub>2</sub>, NO, NO<sub>2</sub>, SCN, OCN, NCS, NCO, SOMe, SO<sub>2</sub>Me;

X is selected from the group consisting of O, S, NH, CH<sub>2</sub>, CHF, CF<sub>2</sub>;

Y is selected from the group consisting of CH<sub>2</sub>, NH, NOH, NMe, NEt, NOME, CHF, CF<sub>2</sub>;

Z is selected from the group consisting of H, OH, OMe, SH, SMe, F, Cl, Br, I, NH<sub>2</sub>, NHMe;

B is a base selected from the group consisting of




R<sup>2</sup> is selected from the group consisting of O, S, NH, NR;



[illegible]


5 consisting of H, OH, OMe, SH, SMe, F, Cl, Br, I, NH<sub>2</sub>, NHMe, NMe<sub>2</sub>;

$R$  is independently selected from the group consisting of







or a pharmaceutically acceptable salt or prodrug thereof, in combination with a pharmaceutically acceptable carrier.

20. The composition of Claim 19, wherein the compound is selected from the group consisting of



21. A pharmaceutical composition for the treatment of HDV comprising an anti-HDV agent and an effective amount of a compound selected from the group consisting of formulas [I]-[IV] below and mixtures of two or more thereof:



wherein:

E is selected from the group consisting of H, OH, OMe, SH, SMe, NH<sub>2</sub>, NHMe, N<sub>3</sub>, F, Cl, Br, CO<sub>2</sub>H, CO<sub>2</sub>-alkyl, OPh, OPhNO<sub>2</sub>, NO, NO<sub>2</sub>, SCN, OCN, NCS, NCO, SOMe, SO<sub>2</sub>Me;



X is selected from the group consisting of O, S, NH, CH<sub>2</sub>, CHF, CF<sub>2</sub>;

Y is selected from the group consisting of CH<sub>2</sub>, NH, NOH, NMe, NEt, NOMe, CHF, CF<sub>2</sub>;

Z is selected from the group consisting of H, OH, OMe, SH, SMe, F, Cl, Br, I, NH<sub>2</sub>,

NHMe;

B is a base selected from the group consisting of




R<sup>2</sup> is selected from the group consisting of O, S, NH, NR;

R<sup>5</sup> is selected from the group consisting of H, branched or unbranched lower alkyl having from 1-5 carbon atoms, F, Cl, Br, I, CH=CH<sub>2</sub>, CH=CHBr, Ph, Ac, OMe, OPh, NO, NO<sub>2</sub>, NH<sub>2</sub>, NHR;


R<sup>6</sup> and R<sup>8</sup> are the same or different and are independently selected from the group consisting of H, OH, OMe, SH, SMe, F, Cl, Br, I, NH<sub>2</sub>, NHMe, NMe<sub>2</sub>;

R is independently selected from the group consisting of







or a pharmaceutically acceptable salt or prodrug thereof, in combination with a pharmaceutically acceptable carrier.

	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	2031	2032	2033	2034	2035	2036	2037	2038	2039	2040	2041	2042	2043	2044	2045	2046	2047	2048	2049	2050	2051	2052	2053	2054	2055	2056	2057	2058	2059	2060	2061	2062	2063	2064	2065	2066	2067	2068	2069	2070	2071	2072	2073	2074	2075	2076	2077	2078	2079	2080	2081	2082	2083	2084	2085	2086	2087	2088	2089	2090	2091	2092	2093	2094	2095	2096	2097	2098	2099	2100	2101	2102	2103	2104	2105	2106	2107	2108	2109	2110	2111	2112	2113	2114	2115	2116	2117	2118	2119	2120	2121	2122	2123	2124	2125	2126	2127	2128	2129	2130	2131	2132	2133	2134	2135	2136	2137	2138	2139	2140	2141	2142	2143	2144	2145	2146	2147	2148	2149	2150	2151	2152	2153	2154	2155	2156	2157	2158	2159	2160	2161	2162	2163	2164	2165	2166	2167	2168	2169	2170	2171	2172	2173	2174	2175	2176	2177	2178	2179	2180	2181	2182	2183	2184	2185	2186	2187	2188	2189	2190	2191	2192	2193	2194	2195	2196	2197	2198	2199	2200	2201	2202	2203	2204	2205	2206	2207	2208	2209	2210	2211	2212	2213	2214	2215	2216	2217	2218	2219	2220	2221	2222	2223	2224	2225	2226	2227	2228	2229	2230	2231	2232	2233	2234	2235	2236	2237	2238	2239	2240	2241	2242	2243	2244	2245	2246	2247	2248	2249	2250	2251	2252	2253	2254	2255	2256	2257	2258	2259	2260	2261	2262	2263	2264	2265	2266	2267	2268	2269	2270	2271	2272	2273	2274	2275	2276	2277	2278	2279	2280	2281	2282	2283	2284	2285	2286	2287	2288	2289	2290	2291	2292	2293	2294	2295	2296	2297	2298	2299	2300	2301	2302	2303	2304	2305	2306	2307	2308	2309	2310	2311	2312	2313	2314	2315	2316	2317	2318	2319	2320	2321	2322	2323	2324	2325	2326	2327	2328	2329	2330	2331	2332	2333	2334	2335	2336	2337	2338	2339	2340	2341	2342	2343	2344	2345	2346	2347	2348	2349	2350	2351	2352	2353	2354	2355	2356	2357	2358	2359	2360	2361	2362	2363	2364	2365	2366	2367	2368	2369	2370	2371	2372	2373	2374	2375	2376	2377	2378	2379	2380	2381	2382	2383	2384	2385	2386	2387	2388	2389	2390	2391	2392	2393	2394	2395	2396	2397	2398	2399	2400	2401	2402	2403	2404	2405	2406	2407	2408	2409	2410	2411	2412	2413	2414	2415	2416	2417	2418	2419	2420	2421	2422	2423	2424	2425	2426	2427	2428	2429	2430	2431	2432	2
--	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	---



23. A method for the treatment of a proliferative disorder comprising administering an effective amount of a compound selected from the group consisting of formulas [I]-[IV] below and mixtures of two or more thereof:



wherein:



E is selected from the group consisting of H, OH, OMe, SH, SMe, NH<sub>2</sub>, NHMe, N<sub>3</sub>, F, Cl, Br, CO<sub>2</sub>H, CO<sub>2</sub>-alkyl, OPh, OPhNO<sub>2</sub>, NO, NO<sub>2</sub>, SCN, OCN, NCS, NCO, SMe, SO<sub>2</sub>Me;

X is selected from the group consisting of O, S, NH, CH<sub>2</sub>, CHF, CF<sub>2</sub>;

Y is selected from the group consisting of CH<sub>2</sub>, NH, NOH, NMe, NEt, NOME, CHF, CF<sub>2</sub>;

Z is selected from the group consisting of H, OH, OMe, SH, SMe, F, Cl, Br, I, NH<sub>2</sub>, NHMe;

B is a base selected from the group consisting of




R<sup>2</sup> is selected from the group consisting of O, S, NH, NR;

R<sup>5</sup> is selected from the group consisting of H, branched or unbranched lower alkyl having from 1-5 carbon atoms, F, Cl, Br, I, CH=CH<sub>2</sub>, CH=CHBr, Ph, Ac, OMe, OPh, NO, NO<sub>2</sub>, NH<sub>2</sub>, NHR;

R<sup>6</sup> and R<sup>8</sup> are the same or different and are independently selected from the group consisting of H, OH, OMe, SH, SMe, F, Cl, Br, I, NH<sub>2</sub>, NHMe, NMe<sub>2</sub>;


R is independently selected from the group consisting of



or a pharmaceutically acceptable salt or prodrug thereof, in combination with a pharmaceutically acceptable carrier.

24. The method of Claim 23, wherein the proliferative disorder is selected from the group consisting of cancer of the stomach, kidney, colon, rectal, liver, pancreatic, lung, breast, cervix uteri, corpus uteri, ovary, prostate, testis, bladder, renal, brain/CNS, head and neck, throat, Hodgkin's disease, non-Hodgkin's leukemia, multiple myeloma leukemias, skin melanoma, acute lymphocytic leukemia, acute myelogenous leukemia, small cell lung cancer, choriocarcinoma, rhabdomyosarcoma, neuroblastoma, mouth/pharynx, esophagus, larynx, melanoma, and lymphoma.


25. The method of Claim 23, wherein the compound is selected from the group consisting of





26. A process for stereospecifically preparing a 5'-modified pyrimidine  $\beta$ -nucleoside comprising

- a. applying the Mitsunobu reaction to a chiral compound of the formula



- b. selectively protecting the 3'- $\beta$ -position of the resulting nucleoside of step (a) with a benzoyl protecting group or an acid labile protecting group;

- c. subjecting the resulting 3'- $\beta$ -protected anhydro derivative of step (b) to mild alkaline hydrolysis, followed by phosphorylating the ring-opened, 3'- $\beta$ -protected product with a phosphorylating agent;


- d. saponification of the benzoyl group of the resulting product of step (c) to give the desired  $\beta$ -nucleoside 5'-phosphate; and

- e. optionally oxidizing the 5'-phosphate to obtain the 5'-phosphite.

27. The process of Claim 26, wherein the acid labile agent is selected from the group consisting of tetrahydropyranyl (THP), a trityl group, or dimethyl-*t*-butylsilyl (DBMS).

28. A process for stereospecifically preparing 5'-phosphorylated pyrimidine  $\alpha$ -nucleosides comprising

- a. applying the Mitsunobu reaction to a chiral compound of the formula




- b. directly phosphorylating the resulting anhydro-derivative of step (a) to yield the  $\alpha$ -nucleoside 5'-phosphate; and

- c. optionally oxidizing the resulting 5'-phosphate to obtain the 5'-phosphite.

29. A process for stereospecifically preparing 5'-phosphorylated pyrimidine  $\alpha$ -nucleosides comprising

- a. treating a compound of the formula



with CBr<sub>4</sub>-Ph<sub>3</sub>P to yield a dibromide;

- b. treating the dibromide with a non-nucleophilic base in an inert solvent to form an anhydro-nucleoside; and

c. treating the anhydro-nucleoside with tribenzylphosphite, followed by hydrogenation to yield the  $\alpha$ -nucleoside 5'-phosphorylated product.

30. A process for preparing purine  $\beta$ -nucleoside 5'-phosphates comprising

- a. converting a purine nucleoside into an 8-thio-derivative by brominating the nucleoside and then treating with thiourea;

b. applying the Mitsunobu reaction to the 8-thio-derivative to yield the S-anhydro product having a  $\beta$ -3'-hydroxy group;

- c. protecting the  $\beta$ -3'-hydroxy group with a base stable protecting group yielding a protected S-anhydro-nucleoside;

d. oxidizing the protected S-anhydro-nucleoside with m-chloroperbenzoic acid;

e. achieving Pummerer rearrangement of the product of step (d) using benzoic anhydride;

f. desulfurizing the product of step (e) with Raney nickel, followed by de-O-benzoylation in base to yield the protected  $\beta$ -3'-nucleoside;

g. phosphorylating the protected  $\beta$ -3'-nucleoside to obtain the  $\beta$ -5'-phosphorylated product; and

h. optionally oxidizing the product of step (g) to obtain the  $\beta$ -5'-phosphite nucleoside.

31. A process for preparing purine  $\alpha$ -nucleoside 5'-phosphates comprising

- a. converting a purine nucleoside into an 8-thio-derivative by brominating the nucleoside and then treating with thiourea;

b. applying the Mitsunobu reaction to the 8-thio-derivative to yield an S-anhydro nucleoside product;

c. phosphorylating the S-anhydro nucleoside to yield a protected S-anhydro nucleoside;

d. oxidizing the protected S-anhydro-nucleoside with m-chloroperbenzoic acid;

e. achieving Pummerer rearrangement of the product of step (d) using benzoic anhydride;

f. desulfurizing the product of step (e) with Raney nickel, followed by de-O-benzoylation in base to yield a phosphorylated  $\beta$ -3'-nucleoside;

g. optionally oxidizing the product of step (f) to obtain the  $\beta$ -3'-phosphite nucleoside.

32. A process for preparing nucleoside 5'-di- and triphosphates comprising

a. addition of sulfur to methylenediphosphite tetrachloride to produce di- or trithiomethylenebis(phosphonate) tetrachloride;

b. hydrolyzing the di- or trithiomethylenebis(phosphonate) tetrachloride to obtain the corresponding phosphonate;

c. treating the resulting compound of step (b) under Yoshikawa-Roeschler's conditions to yield the corresponding P<sup>1</sup>-nucleoside-5'-yl-methylenebis-(thiophosphonate);

d. S-alkylating and treating the resulting compound of step (c) with fluoride (Bu<sub>4</sub>NF, DAST) to yield fluorine substituted 3'- $\beta$ -protected nucleoside;

e. deprotecting the 3'- $\beta$ -nucleoside to obtain the 5'-di- or 5'-triphosphate  $\beta$ -nucleoside.